

		CD10	BCL6	PD-1	CXCR5	ICOS	TCF1	TBX21	CXCR3	GATA3	CCR4
CD10	Pearson's r		.339**	.372**	.260*	0.231	0.115	-0.075	0.097	-0.245	0.145
	P value		0.007	0.003	0.041	0.071	0.43	0.593	0.486	0.074	0.297
BCL6	Pearson's r			0.1	0.243	.332**	0.132	0.097	0.217	-0.235	-0.137
	P value			0.438	0.057	0.008	0.367	0.491	0.114	0.087	0.322
PD-1	Pearson's r				.575**	0.173	0.133	0.146	0.084	-.314*	0.139
	P value				0	0.178	0.364	0.296	0.547	0.021	0.317
CXCR5	Pearson's r					.317*	.317*	0.092	0.216	-.339*	0.005
	P value					0.012	0.026	0.511	0.117	0.012	0.972
ICOS	Pearson's r						.386**	-0.123	0.09	-0.059	-0.087
	P value						0.006	0.379	0.516	0.674	0.531
TCF1	Pearson's r							0.084	.290*	0.132	0.036
	P value							0.503	0.018	0.292	0.775
TBX21	Pearson's r								0.063	0.017	-0.044
	P value								0.589	0.887	0.712
CXCR3	Pearson's r									-0.168	-.231*
	P value									0.147	0.046
GATA3	Pearson's r										.354**
	P value										0.002

* $P < 0.05$, ** $P < 0.01$

Supplementary Fig. S2. Correlation between Tfh, TCF-1, and Th1- and Th2-like markers. CXCR5, C-X-C motif chemokine receptor 5; GATA3, GATA binding protein 3; ICOS, inducible T cell costimulator; PD-1, programmed death-1; TBX21, T-box transcription factor 21; TCF1, T-cell factor-1; Tfh, follicular helper T-cell